

Non-linear realization of Poincaré invariance in the graph-representation of extremal hypersurfaces

Jens Hoppe *

Department of Mathematics,
Royal Institute of Technology,
KTH, 100 44 Stockholm,
Sweden

Abstract

In the Born-Infeld 'harmonic gauge' description of M-branes moving in $\mathbb{R}^{M+1}$ the underlying $M + 2$ dimensional Poincaré - invariance gives rise to an interesting system of conservation laws showing signs of integrability.

Varying

$$S[z] = - \int d^M x dt \sqrt{1 - \dot{z}^2 + (\vec{\nabla} z)^2} \quad (1)$$

yields the quasi-linear PDE

$$(1 - z^\alpha z_\alpha) \square z + z^\alpha z^\beta z_{\alpha\beta} = 0 \quad (2)$$

for $z(t, \vec{x})$, with $z^\alpha := \eta^{\alpha\beta} z_\beta = \eta^{\alpha\beta} \frac{\partial z}{\partial x^\beta}$, $(\alpha, \beta = 0, 1, \dots, M)$, $\eta^{\alpha\beta} = \text{diag}(1, -1, \dots, -1)$, $z_{\alpha\beta} = \frac{\partial^2 z}{\partial x^\alpha \partial x^\beta}$, $\square z = \eta^{\alpha\beta} z_{\alpha\beta}$.

At first sight, (2) appears to be a scalar field equation in $M + 1$ - dimensional Minkowski space, but $M + 2$ additional conservation laws follow when

*e-mail: hoppe@math.kth.se

realizing that z is most naturally interpreted as a (for notational convenience, the last) component of a $(M+2)$ - vector

$$x^\mu = \begin{pmatrix} t \\ \vec{x} \\ z(t, \vec{x}) \end{pmatrix} \quad (3)$$

in $\mathbb{R}^{1,M+1}$ (cp. e.g. [1]). The corresponding world-volume $\mathcal{M}$, described by (3)/(2), can easily be shown to have vanishing mean curvature. The $M+2$ -dimensional Poincaré - symmetry (non-linearly realized, as $x^{M+1} = z(t, \vec{x})$) is expressed by

$$\partial_\alpha \mathcal{H}^{\alpha\mu} = 0 \quad (\mu = 0, 1, \dots, M+1) \quad (4)$$

$$\partial_\alpha (x^\mu \mathcal{H}^{\nu\alpha} - x^\nu \mathcal{H}^{\mu\alpha}) = 0 \quad (5)$$

$$\mathcal{H}^{\alpha\beta} := \frac{z^\alpha z^\beta}{\sqrt{1 - z^\gamma z_\gamma}} + \eta^{\alpha\beta} \sqrt{1 - z^\gamma z_\gamma} \quad (6)$$

$$\mathcal{H}^{\alpha, M+1} := \frac{z^\alpha}{\sqrt{1 - z^\gamma z_\gamma}} = \mathcal{H}^{M+1, \alpha} \quad (7)$$

(generalising to arbitrary co-dimension is straightforward).

Note in particular that

$$z_\alpha \mathcal{H}^{\alpha\beta} = \mathcal{H}^{\alpha, M+1} \quad (8)$$

Let me make a few comments how one may be able to *use* (4) and (5).

First of all note the monotonicity of

$$\int x^\mu \mathcal{H}^{00} = C^\mu + t P^\mu \quad (9)$$

(C^μ and $P^\mu = \int \mathcal{H}^{\mu 0}$ being constants, depending only on the initial conditions); and the explicit (polynomial; here linear) appearance of t .

Secondly, one could hope that the conservation laws, in particular the $\mu \neq M+1$ part of (4) (containing neither x^α nor z !), thought of as a *first-order* system linear in the derivatives of the $M+1$ -vector

$$z^\alpha = \begin{pmatrix} \dot{z} \\ -\vec{\nabla} z \end{pmatrix}, \quad (10)$$

resp. $u^\alpha = \frac{z^\alpha}{\sqrt{1 - z^\gamma z_\gamma}}$, resp. $\mathcal{H}^{\alpha 0}$ (or other suitably chosen coordinates; e.g. in order to apply the method of characteristics) are *integrable*.

Acknowledgments I would like to thank M. Bordemann, B. Gustafsson, E. Langmann, J. Reinhardt, M. Trzetrzelewski and A. Zheltukhin for valuable discussions.

References

- [1] M. Bordemann, J. Hoppe, PLB 325 (1994) 359-365,